

Flea Presence and Abundance Are Not Predictors of *Bartonella tribocorum* Carriage in Norway Rats (*Rattus norvegicus*) from an Underserved Neighborhood of Vancouver, Canada

AU1 ► Chelsea G. Himsworth,¹⁻³ Kaylee A. Byers,^{1,4,5} Tammi Whelan,¹ Ying Bai,⁶ and Michael Y. Kosoy⁷

AU3 ► Abstract

Urban Norway rats (*Rattus norvegicus*) carry pathogenic *Bartonella* spp. that are transmitted among rats and from rats to people through arthropod vectors, particularly fleas. There is marked temporospatial variation in *Bartonella* spp. carriage among Norway rats in Vancouver, Canada, and we investigated whether this variation is associated with flea presence or abundance. *Bartonella tribocorum* was isolated from 96/370 (35%) rats and 211 (57%) rats had fleas with an average of one flea per rat. All fleas were identified as *Nosopsyllus fasciatus*. There was no significant relationship between *B. tribocorum* carriage and flea presence or abundance, suggesting that, in contrast to other rat-associated zoonoses transmitted by fleas (e.g., *Yersinia pestis*) flea indices may not be informative for understanding the ecology of *Bartonella* spp. in rats, particularly for *N. fasciatus*.

Keywords: *Bartonella* spp., fleas, rats, *Rattus norvegicus*, *Nosopsyllus fasciatus*, urban

Introduction

RATS (*RATTUS* spp.) ARE the reservoir for a variety of zoonotic pathogens (Himsworth et al. 2013), including *Bartonella* spp.—a genus of vector-borne intracellular bacteria that infect endothelial cells and erythrocytes (Kosoy and Bai 2019). At least 22 *Bartonella* spp. have been found in 98 rodent species, with *Bartonella tribocorum*, *Bartonella elizabethae*, *Bartonella rattimalssiliensis*, and *Bartonella queenslandensis* being associated with rats (Kosoy and Bai 2019). *Bartonella* spp. are transmitted among rats and from rats to people by hematophagous arthropods, particularly fleas (Gutierrez et al. 2015). In rats, *Bartonella* spp. cause a persistent asymptomatic bacteremia (Gutierrez et al. 2015). In people, infection with rat-associated *Bartonella* spp. can cause septicemia with a range of clinical presentations (Kosoy and Bai 2019).

Understanding the ecology of zoonotic pathogens in the reservoir host is critical for monitoring and mitigating the

risk of spillover from rats to people. The ecology of *Bartonella* spp. in rodents is dependent on a complex set of interactions among the *Bartonella* spp., the rodent host, the environment, and the vector(s) (Gutierrez et al. 2015). In other rodent species, it is thought that flea abundance may be an important determinant of infection (Gutierrez et al. 2015), yet the role of fleas in the ecology of *Bartonella* spp. in urban rats is unclear.

A previous study of urban Norway rats (*Rattus norvegicus*) from Vancouver, Canada, found significant unexplained temporospatial variation in *B. tribocorum* carriage (Himsworth et al. 2015). The objective of this study is to determine whether this variation could be attributed to the presence and abundance of fleas infesting the rats under study.

Materials and Methods

For the aforementioned previous study (Himsworth et al. 2015), rats were trapped in 43 contiguous city blocks over the

AU2 ►

¹Canadian Wildlife Health Cooperative—British Columbia, Abbotsford, Canada.

²School of Population and Public Health, University of British Columbia, Vancouver, Canada.

³Animal Health Centre, British Columbia Ministry of Agriculture, Abbotsford, Canada.

⁴Department of Interdisciplinary Studies, University of British Columbia, Vancouver, Canada.

⁵Biodiversity Research Centre, University of British Columbia, Vancouver, Canada.

⁶Division of Vector-Borne Diseases, Centers for Disease Control and Prevention, Fort Collins, Colorado, USA.

⁷KB One Health, Fort Collins, Colorado, USA.

TABLE 1. DISTRIBUTION OF EXPLANATORY VARIABLES AND ASSOCIATIONS WITH *BARTONELLA TRIBOCORUM* STATUS AMONG NORWAY RATS


Category	Subcategory	Total (%) (n=370)	<i>Bartonella tribocorum</i> status		Bivariable GLMM ^a	
			Positive (%) (n=96)	Negative (%) (n=274)	OR (95% CI)	p
Season	Fall	150 (40.5)	74 (77.1)	76 (27.7)	Ref	
	Winter	77 (20.8)	5 (6.5)	72 (26.3)	0.08 (0.02–0.45)	<0.001
	Spring	103 (27.8)	12 (12.5)	91 (33.2)	0.09 (0.02–0.38)	<0.01
	Summer	40 (10.8)	5 (5.2)	35 (12.8)	0.07 (0.01–0.59)	0.01
Maturity	Immature	89 (24.1)	9 (9.4)	80 (29.2)	Ref	
	Mature	281 (75.9)	87 (90.6)	194 (70.8)	2.45 (1.07–5.64)	0.03
Flea presence	No	159 (43.0)	35 (36.5)	124 (45.3)	Ref	
	Yes	211 (57.0)	61 (63.5)	150 (54.7)	1.17 (0.65–2.11)	0.59
Flea abundance	0	159 (43.0)	35 (36.5)	124 (45.3)	Ref	
	1–3	167 (45.1)	45 (46.9)	122 (44.5)	1.10 (0.60–2.04)	0.76
	>3	44 (11.9)	16 (16.7)	28 (10.2)	1.48 (0.61–3.61)	0.39
Flea index	≤0.65	151 (40.8)	53 (55.2)	98 (35.8)	Ref	
	0.65–1.3	107 (28.9)	26 (27.1)	81 (29.6)	0.39 (0.05–3.25)	0.39
	>1.3	112 (30.3)	17 (17.7)	95 (34.7)	0.21 (0.03–1.47)	0.12

AU6 ▶ ^aGLMM controlling for clustering by city block in which the trap was trapped. CI, confidence interval; GLMM, generalized linear mixed model; OR, odds ratio.

course of 1 year. Blood was collected through intracardiac puncture under isoflurane anesthesia before pentobarbital euthanasia and, subsequently, fleas were collected by combing the fur. For each rat, the date and location of trapping, species, and maturity were recorded (animals were considered sexually mature if they had scrotal testes or a

perforate vagina). Blood clots underwent *Bartonella* culture and isolates were identified by *gltA* sequencing, while fleas were stored at –80°C.

For this study, collected fleas were examined using a compound microscope (40×) for enumeration and identification to species (Holland 1985). Bivariate and multivariate



4C ▶

FIG. 1. Distribution of *Bartonella tribocorum*-positive Norway rats (*Rattus norvegicus*) and flea index by city block.

generalized linear mixed models (GLMMs) controlling for clustering by city block of origin were used to assess whether flea indices are associated with *Bartonella* spp. carriage in rats. The unit of analysis was the rat and the outcome variable was *Bartonella* spp. carriage (positive vs. negative based on culture and sequencing of blood clots). Explanatory variables included for consideration included flea presence (0 flea/rat vs. ≥ 1 flea/rat), flea abundance (number of fleas per rat), and block-level flea index (average number of fleas per rat for the city block in which the rat was trapped). Sexual maturity (immature vs. mature) and season (fall: September–November; winter: December–February; spring: March–May; summer: June–August) were forced into the model as these variables had been identified as predictors of *Bartonella* spp. carriage in previous models of these data (Himsworth et al., 2015). Variables that were significantly associated with *Bartonella* spp. carriage at an alpha level of ≤ 0.10 in a bivariable GLMM were considered for inclusion in multivariable modelling. The Kruskal–Wallis test was used to assess seasonal differences in flea abundance and flea index. All statistical analyses were conducted using R (R Development Core Team, Vienna, Austria). The trap location and number of *B. tribocorum*-positive and -negative rats as well as the block-level flea index were mapped using ArcGIS 10.0 (ESRI, Redlands, USA).

This study was approved by the University of British Columbia's Animal Care Committee (A11-0087).

Results

A total of 370 Norway rats were included in this study of which 96 (35%) were positive for *B. tribocorum*. Fleas were present on 211 (57%) of rats with a median of one flea per rat (range 0–14). All fleas were identified as *Nosopsyllus fasciatus*. The median flea index for a city block was 0.93 (range 0–5.6). Both flea abundance and flea index demonstrated significant ($p < 0.01$) seasonal variation—being highest in the spring, lowest in the winter, and intermediate in the summer and fall. There was no significant relationship between *B. tribocorum* carriage and flea presence, flea abundance, or flea index (Table 1). Mapping also showed that block-level flea index was not clearly correlated with the number of rats carrying *B. tribocorum* in that block (Fig. 1). The odds of *B. tribocorum* carriage were increased in mature animals and decreased for rats trapped in the spring, summer, and winter compared with the fall (Table 1), consistent with previous models of these data (Himsworth et al. 2015).

Discussion

Although many studies have examined the presence and diversity of *Bartonella* spp. in urban rats and their fleas (Kosoy and Bai 2019), far fewer have sought to determine how flea ecology correlates with pathogen prevalence and distribution in the rat host. Peterson et al. (2017) found that Norway rats that had fleas (*Xenopsylla cheopis*) were more likely to be infected with *Bartonella* spp. than those that did not. However, we did not identify an association between *B. tribocorum* carriage in Norway rats and flea presence, flea abundance, or block-level flea index.

Given that *Bartonella* spp. can cause a persistent bacteremia the fleas present on a rat at the time of trapping are unlikely to be those present at the time of infection. In addition, for *N. fasciatus*, a species that spends more time in the rat's nest compared with *X. cheopis*, the number of fleas on the rat at any given time may not reflect the number of fleas to which that rat has been exposed (Bitam et al. 2010). There has only been a single study of *N. fasciatus* as a vector for *Bartonella* spp. in rodents (specifically, *Rattus surifer* in Thailand (Parola et al. 2003). This paucity of information is unexpected as *N. fasciatus* is reported to be common on commensal Norway rats in temperate regions (Bitam et al. 2010).

It is interesting to note that rat-level flea abundance and block-level flea index were highest in the spring, whereas *Bartonella* spp. was most prevalent in the fall. *Bartonella* spp. carriage and flea abundance/index could, therefore, be linked, but temporally offset in such a manner that would not be revealed in a GLMM. Alternatively, it may be the case that there are seasonal variations in vector activity/competence and/or host–vector interactions. A limitation of this study is the fact that the fleas were not tested for *Bartonella* spp. and future studies should seek to clarify the relationship between *Bartonella* spp. carriage in rats and the fleas that infest them.

Conclusion

This study suggests that, in contrast to other rat-associated zoonoses transmitted by fleas, such as *Yersinia pestis* (Dennis et al. 1999), flea indices may not be informative for understanding the ecology of *Bartonella* spp. in rats, particularly for *N. fasciatus*.

Acknowledgment

Flea identification confirmed was performed by Dr. Terry Galloway, University of Manitoba.

Author Disclosure Statement

No conflicting financial interests exist.

Funding Information

◀ AU4

References

- Bitam I, Dittmar K, Parola P, Whiting MF, et al. Fleas and flea-borne diseases. *Int J Infect Dis* 2010;14:e667–e676.
- Dennis DT, Gage KL, Gratz NG, Poland JD, et al. *Plague Manual: Epidemiology, Distribution, Surveillance and Control*. World Health Organization, 1999.
- Gutierrez R, Krasnov B, Morick D, Gottlieb Y, et al. *Bartonella* infection in rodents and their flea ectoparasites: An overview. *Vector Borne Zoonotic Dis* 2015;15:27–39.
- Himsworth CG, Bai Y, Kosoy MY, Wood H, et al. An investigation of *Bartonella* spp., *Rickettsia typhi*, and Seoul hantavirus in rats (*Rattus* spp.) from an inner-city neighborhood of Vancouver, Canada: Is pathogen presence a reflection of global and local rat population structure? *Vector Borne Zoonotic Dis* 2015;15:21–26.

◀ AU5

- Himsworth CG, Parsons KL, Jardine C, Patrick DM. Rats, cities, people, and pathogens: A systematic review and narrative synthesis of literature regarding the ecology of rat-associated zoonoses in urban centers. *Vector Borne Zoonotic Dis* 2013; 13:349–359.
- Holland GP. The Fleas of Canada, Alaska, and Greenland (Siphonaptera). *Memoirs Entomol Soc Canada* 1985;117 (S130): 3–632.
- Kosoy M, Bai Y. *Bartonella* bacteria in urban rats: A movement from the jungles of Southeast Asia to metropolises around the globe. *Front Ecol Evol* 2019;7:88.
- Parola P, Sanogo OY, Lerdthusneem K, Zeaiter Z, et al. Identification of *Rickettsia* spp. and *Bartonella* spp. in fleas from the Thai-Myanmar border. *Ann N Y Acad Sci* 2003; 990:173–181.
- Peterson AC, Ghersi BM, Alda F, Firth C, et al. Rodent-borne *Bartonella* infection varies according to host species within and among cities. *EcoHealth* 2017;14:771–782.


Address correspondence to:
Chelsea G. Himsworth
Animal Health Centre
British Columbia Ministry of Agriculture
1767 Angus Campbell Road
Abbotsford
British Columbia V3G 2M3
Canada


E-mail: chelsea.himsworth@gov.bc.ca

AUTHOR QUERY FOR VBZ-2020-2665-VER9-HIMSWORTH_1P

AU1: Please identify (highlight or circle) all authors' surnames for accurate indexing citations.

AU2: Please check the correctness of authors' affiliations.

AU3: Include IRB approval or waiver statement in the Materials and Methods section. The Clinical Trial Registration number, if applicable, should be included at the end of the abstract. 

AU4: Please provide funding information. Include the full name (no abbreviations) of funding agency(ies)/institution(s) and grant numbers. Ensure the accuracy of this information. If no funding was received, please clearly state as such. 

AU5: In Ref. "Dennis et al. (1999)," please mention the publisher's location. 

AU6: Please check the footnote in Table 1: is it "trap was trapped" or "rat was trapped"? 